



Serbia

Air pollution was the 7th leading risk factor for premature death in Serbia in 2019, accounting for nearly 11% of all deaths. Considered separately, ambient particulate matter (PM_{2.5}) ranked as the 7th leading risk factor, and household air pollution (HAP) ranked 16th. Ozone was not in the top 20 risk factors.

Key Statistics at a Glance


<p>Nearly 13,000 deaths due to air pollution in 2019.</p> <p>Less than 10% of infant deaths attributable to air pollution.</p>	 <p>25 µg/m³ population-weighted annual average PM_{2.5} concentration.*</p> <p>More than 10,500 deaths attributable to exposure to outdoor PM_{2.5}.</p>	 <p>24% of the population used solid fuels for cooking.</p> <p>Nearly 2,000 deaths attributable to exposure to household air pollution.</p>
--	---	--

Key Exposure Facts

100% of Serbia's population lives in areas where PM_{2.5} levels are above the WHO guideline for healthy air (10 µg/m³).**

- Between 2010 and 2019, exposures to PM_{2.5}, household air pollution, and ozone all declined.
- There are less than 10 stations reporting PM_{2.5} concentrations in Serbia.***
- Among the 29 countries in the Central Europe, Eastern Europe, and Central Asia region, Serbia ranks 9th in PM_{2.5} exposure.

How Have Pollutant Exposures Changed Between 2010 and 2019?



* Please note that PM_{2.5} concentrations reported here are estimated using a combination of satellite data, ground air quality monitoring data, and chemical transport models. These estimates can be more uncertain in regions where ground monitoring data are limited or not available. In Serbia, the best estimate of the annual average exposure is 25 µg/m³, but it may range from 23 µg/m³ to 28 µg/m³.

** WHO provides an Air Quality Guideline of 10 µg/m³ for PM_{2.5} to minimize health risks to populations, as well as three interim targets (35 µg/m³, 25 µg/m³, and 15 µg/m³) as incremental steps toward the progressive reduction of air pollution.


*** Based on data from OpenAQ.

STATE OF GLOBAL AIR /2020

Air Pollution Accounts for a Substantial Percentage of Global Deaths from Specific Causes.

Air pollution exposures, including exposure to outdoor PM_{2.5} and HAP, have been linked to increased hospitalizations, disability, and early death from respiratory diseases, heart disease, stroke, lung cancer, and diabetes, as well as communicable diseases like pneumonia. Exposure to ozone is linked to chronic obstructive pulmonary disease (COPD), and in children, especially those under the age of 5, increases susceptibility to lower respiratory tract infections. Exposure to PM_{2.5} also puts mothers at risk of delivering babies too early and smaller than normal, and such babies are more susceptible to dying from a range of diseases.

Percentage of Deaths (by Cause) Attributed to Air Pollution in Serbia in 2019



Key Health Facts

- Air pollution is the 7th leading risk factor for premature death in Serbia. Leading causes of death in Serbia include ischemic heart disease, ischemic stroke, lung cancer, intracerebral hemorrhage, and colorectal cancer, while leading risk factors include high blood pressure, dietary risks, tobacco, high blood sugar, and high BMI.
- There are 83 deaths per 100,000 people attributable to air pollution in Serbia compared with 86 deaths globally, adjusted for differences in age.
- 5% of total air-pollution-attributable deaths in Serbia are in children under 5, and 11% are in people over 70.

FOR MORE INFORMATION:

For the full report and additional data, please visit www.stateofglobalair.org.

ADDITIONAL RESOURCES:

For open-access, real-time air quality data, visit OpenAQ



For more details, please visit www.stateofglobalair.org
Contact us contactsoga@healtheffects.org



IHME



The State of Global Air website is a collaboration between the Health Effects Institute and the Institute for Health Metrics and Evaluation, with expert input from the University of British Columbia.